

CONGRUENCES ON THE MONOID OF MONOTONE INJECTIVE PARTIAL SELFMAPS OF $L_n \times_{\text{lex}} \mathbb{Z}$ WITH CO-FINITE DOMAINS AND IMAGES

OLEG GUTIK AND INNA POZDNIAKOVA

ABSTRACT. We study congruences on the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ of monotone injective partial selfmaps of the set of $L_n \times_{\text{lex}} \mathbb{Z}$ having co-finite domains and images, where $L_n \times_{\text{lex}} \mathbb{Z}$ is the lexicographic product of n -elements chain and the set of integers with the usual linear order. The structure of the sublattice of congruences on $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ which contained in the least group congruence is described.

We follow the terminology of [5, 6] and [7]. We shall denote the additive group of integers by $\mathbb{Z}(+)$.

An algebraic semigroup S is called *inverse* if for any element $x \in S$ there exists a unique $x^{-1} \in S$ such that $xx^{-1}x = x$ and $x^{-1}xx^{-1} = x^{-1}$. The element x^{-1} is called the *inverse of $x \in S$* . If S is an inverse semigroup, then the function $\text{inv}: S \rightarrow S$ which assigns to every element x of S its inverse element x^{-1} is called an *inversion*.

If $\mathfrak{C}$ is an arbitrary congruence on a semigroup S , then we denote by $\Phi_{\mathfrak{C}}: S \rightarrow S/\mathfrak{C}$ the natural homomorphisms from S onto the quotient semigroup $S/\mathfrak{C}$. A congruence $\mathfrak{C}$ on a semigroup S is called *non-trivial* if $\mathfrak{C}$ is distinct from universal and identity congruences Δ_S on S , and *group* if the quotient semigroup $S/\mathfrak{C}$ is a group. Every inverse semigroup S admits the least (minimum) group congruence σ :

$$a\sigma b \text{ if and only if there exists } e \in E(S) \text{ such that } ae = be$$

(see [7, Lemma III.5.2])

If S is a semigroup, then we shall denote the subset of idempotents of S by $E(S)$. If S is an inverse semigroup, then $E(S)$ is closed under multiplication and we shall refer to $E(S)$ as a *band* (or the *band of S*). If the band $E(S)$ is a non-empty subset of S , then the semigroup operation on S determines the following partial order $\leq$ on $E(S)$: $e \leq f$ if and only if $ef = fe = e$. This order is called the *natural partial order* on $E(S)$. A *semilattice* is a commutative semigroup of idempotents. A semilattice E is called *linearly ordered* or a *chain* if its natural order is a linear order. A *maximal chain* of a semilattice E is a chain which is properly contained in no other chain of E .

If S is a semigroup, then we shall denote the Green relations on S by $\mathcal{R}$, $\mathcal{L}$, $\mathcal{J}$, $\mathcal{D}$ and $\mathcal{H}$ (see [1, Section 2.1]):

$$\begin{aligned} a\mathcal{R}b &\text{ if and only if } aS^1 = bS^1; \\ a\mathcal{L}b &\text{ if and only if } S^1a = S^1b; \\ a\mathcal{J}b &\text{ if and only if } S^1aS^1 = S^1bS^1; \\ \mathcal{D} &= \mathcal{L} \circ \mathcal{R} = \mathcal{R} \circ \mathcal{L}; \\ \mathcal{H} &= \mathcal{L} \cap \mathcal{R}. \end{aligned}$$

A semigroup S is called *simple* if S contains no proper two-sided ideal, i.e., S has a unique $\mathcal{J}$ -class, and *bisimple* if S has a unique $\mathcal{D}$ -class.

If $\alpha: X \rightarrow Y$ is a partial map, then by $\text{dom } \alpha$ and $\text{ran } \alpha$ we denote the domain and the range of α , respectively.

Date: May 26, 2015.

2010 Mathematics Subject Classification. Primary 20M18, 20M20. Secondary 20M05, 20M15.

Key words and phrases. Semigroup of bijective partial transformations, symmetric inverse semigroup, congruence.

Let $\mathcal{I}_\lambda$ denotes the set of all partial one-to-one transformations of an infinite set X of cardinality λ endowed with the following semigroup operation: $x(\alpha\beta) = (x\alpha)\beta$ if $x \in \text{dom}(\alpha\beta) = \{y \in \text{dom } \alpha \mid y\alpha \in \text{dom } \beta\}$, for $\alpha, \beta \in \mathcal{I}_\lambda$. The semigroup $\mathcal{I}_\lambda$ is called the *symmetric inverse semigroup* over the set X (see [1, Section 1.9]). The symmetric inverse semigroup was introduced by Wagner [8] and it plays a major role in the theory of semigroups. An element $\alpha \in \mathcal{I}_\lambda$ is called *cofinite*, if the sets $\lambda \setminus \text{dom } \alpha$ and $\lambda \setminus \text{ran } \alpha$ are finite.

Let $(X, \leq)$ be a partially ordered set. We shall say that a partial map $\alpha: X \rightarrow X$ is *monotone* if $x \leq y$ implies $(x)\alpha \leq (y)\alpha$ for each $x, y \in X$.

Let $\mathbb{Z}$ be the set of integers with the usual linear order " $\leq$ ". For any positive integer n by L_n we denote the set $\{1, \dots, n\}$ with the usual linear order " $\leq$ ". On the Cartesian product $L_n \times \mathbb{Z}$ we define the lexicographic order, i.e.,

$$(i, m) \leq (j, n) \quad \text{if and only if} \quad (i < j) \quad \text{or} \quad (i = j \quad \text{and} \quad m \leq n).$$

Later the set $L_n \times \mathbb{Z}$ with the lexicographic order we denote by $L_n \times_{\text{lex}} \mathbb{Z}$. Also, it is obvious that the set $\mathbb{Z} \times L_n$ with the lexicographic order is order isomorphic to $(\mathbb{Z}, \leq)$.

By $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ we denote a semigroup of injective partial monotone selfmaps of $L_n \times_{\text{lex}} \mathbb{Z}$ with cofinite domains and images. Obviously, $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is an inverse submonoid of the semigroup $\mathcal{I}_\omega$ and $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is a countable semigroup. Also, by $\mathcal{IO}_\infty(\mathbb{Z})$ we denote a semigroup of injective partial monotone selfmaps of $\mathbb{Z}$ with cofinite domains and images.

Furthermore, we shall denote the identity of the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ by $\mathbb{I}$ and the group of units of $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ by $H(\mathbb{I})$.

Gutik and Repovš in [3] showed that the semigroup $\mathcal{I}_\infty^\nearrow(\mathbb{N})$ of partial cofinite monotone injective transformations of the set of positive integers $\mathbb{N}$ has algebraic properties similar to those of the bicyclic semigroup: it is bisimple and all of its non-trivial semigroup homomorphisms are either isomorphisms or group homomorphisms.

In [4] Gutik and Repovš studied the semigroup $\mathcal{I}_\infty^\nearrow(\mathbb{Z})$ of partial cofinite monotone injective transformations of the set of integers $\mathbb{Z}$ and they showed that $\mathcal{I}_\infty^\nearrow(\mathbb{Z})$ is bisimple and all of its non-trivial semigroup homomorphisms are either isomorphisms or group homomorphisms.

In the paper [2] we studied the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. There we described Green's relations on $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$, showed that the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is bisimple and established its projective congruences. Also, there we proved that $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is finitely generated, every automorphism of $\mathcal{IO}_\infty(\mathbb{Z})$ is inner and showed that in the case $n \geq 2$ the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ has non-inner automorphisms. In [2] we proved that for every positive integer n the quotient semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)/\sigma$, where σ is the least group congruence on $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$, is isomorphic to the direct power $(\mathbb{Z}(+))^{2^n}$.

By Proposition 2.3(iv) [2], the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is isomorphic to the direct power $(\mathcal{IO}_\infty(\mathbb{Z}))^n$. Fixing this isomorphism further we shall identify elements of the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ with elements of the direct product $(\mathcal{IO}_\infty(\mathbb{Z}))^n$, i.e., every element α of $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ we present in the form $(\alpha_1, \alpha_2, \dots, \alpha_n)$, where all α_i belongs to $\mathcal{IO}_\infty(\mathbb{Z})$. Later by α_i° we shall denote the element with the form $(\mathbb{I}_1, \dots, \mathbb{I}_{i-1}, \alpha_i, \mathbb{I}_{i+1}, \dots, \mathbb{I}_n)$, where $\mathbb{I}_j$ is the identity of the j -th factor of $(\mathcal{IO}_\infty(\mathbb{Z}))^n$ for all j and $\alpha_i \in (\mathcal{IO}_\infty(\mathbb{Z}))$. It is obvious that for every $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ we have that $\alpha = \alpha_1^\circ \dots \alpha_n^\circ$.

For every $i = 1, \dots, n$ we define a binary relation $\sigma_{[i]}$ on the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ in the following way:

$$\alpha \sigma_{[i]} \beta \quad \text{if and only if there exists an idempotent } \varepsilon \in \mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n) \text{ such that } \alpha \varepsilon_i^\circ = \beta \varepsilon_i^\circ.$$

In [2] we proved that $\sigma_{[i]}$ is a congruence on $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ for every $i = 1, \dots, n$. Also, there is shown that for any subset $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ of distinct integers, the relation $\sigma_{[i_1, \dots, i_k]} = \sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]}$ is a congruence on $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ and is described the properties of the congruence $\sigma_{[i_1, \dots, i_k]}$ (see Propositions 2.11-2.13, 2.15 and 2.18 in [2]). Moreover, $\sigma_{[1, 2, \dots, n]}$ is the least group congruence on the semigroup $\mathcal{IO}_\infty(\mathbb{Z}_{\text{lex}}^n)$.

For every $i = 1, \dots, n$ we define a map $\pi^i: \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n) \rightarrow \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ by the formula $(\alpha)\pi^i = \alpha_i^\circ$, i.e., $(\alpha_1, \dots, \alpha_i, \dots, \alpha_n)\pi^i = (\mathbb{I}_1, \dots, \mathbb{I}_{i-1}, \alpha_i, \mathbb{I}_{i+1}, \dots, \mathbb{I}_n)$. Simple verifications show that the map $\pi^i: \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n) \rightarrow \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is a homomorphism. Let $\pi^{i\sharp}$ be the congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ which is generated by the homomorphism π^i .

Let S be an inverse semigroup. For any congruence ρ on S we define a congruence $\rho_{\min}$ on S as follows:

$$a\rho_{\min}b \quad \text{if and only if} \quad ae = be \quad \text{for some} \quad e \in E(S) \quad \text{and} \quad epa^{-1}apb^{-1}b,$$

(see: [7, Section III.2]). Then Proposition 2.17 of [2] implies that

$$\pi^{i\sharp}_{\min} = \sigma_{[1]} \circ \dots \circ \sigma_{[i-1]} \circ \sigma_{[i+1]} \circ \dots \circ \sigma_{[n]}$$

for every $i = 1, \dots, n$.

This paper is a continuation of [2] and we study congruences on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Here we describe the structure of the sublattice of congruences on $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ which contained in the least group congruence.

For arbitrary elements $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$ of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ we define:

$$\mathbf{D}_{\alpha, \beta} = \{i \in \{1, \dots, n\} \mid \alpha_i \neq \beta_i\}.$$

It is obvious that elements $\alpha, \beta \in \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ are equal if and only if $\mathbf{D}_{\alpha, \beta} = \emptyset$.

Lemma 1. *Let $\mathfrak{C}$ be a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Let α and β be two distinct $\mathfrak{C}$ -equivalent elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Then there exists an element ϖ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbb{I}\mathfrak{C}\varpi$ and $\mathbf{D}_{\mathbb{I}, \varpi} = \mathbf{D}_{\alpha, \beta}$.*

Proof. By Proposition 2.3(iv) from [2] the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is isomorphic to the direct power $(\mathcal{SO}_\infty(\mathbb{Z}))^n$. We denote $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$. Then for every $i \in \mathbf{D}_{\alpha, \beta}$ we have that $\alpha_i \neq \beta_i$.

We fix an arbitrary $i \in \mathbf{D}_{\alpha, \beta}$. Then one of the following cases holds:

- 1) $\alpha_i \mathcal{H} \beta_i$ in $\mathcal{SO}_\infty(\mathbb{Z})$;
- 2) α_i and β_i are not $\mathcal{H}$ -equivalent in $\mathcal{SO}_\infty(\mathbb{Z})$.

Suppose that case 1) holds. By Proposition 2.3 of [4] the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is bisimple and hence by Theorem 2.3 from [1] there exist $\gamma_i, \delta_i \in \mathcal{SO}_\infty(\mathbb{Z})$ such that $\eta_i = \gamma_i \alpha_i \delta_i$ and $\zeta_i = \gamma_i \beta_i \delta_i$ are distinct elements of the group of units of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$. Then we have that $\eta_i^{-1} \eta_i = \eta_i^{-1} \gamma_i \alpha_i \delta_i = \mathbb{I}_i$ is the unit of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ and $\eta_i^{-1} \zeta_i = \eta_i^{-1} \gamma_i \beta_i \delta_i \neq \mathbb{I}_i$. Hence, without loss of generality we can assume that there exist elements γ_i and δ_i of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ such that $\gamma_i \alpha_i \delta_i = \mathbb{I}_i$ is the unit of $\mathcal{SO}_\infty(\mathbb{Z})$ and $\gamma_i \beta_i \delta_i \neq \mathbb{I}_i$.

Suppose that the elements α_i and β_i are not $\mathcal{H}$ -equivalent in $\mathcal{SO}_\infty(\mathbb{Z})$. Then by Proposition 2.1(vii) of [4] we have that at least one of the following conditions holds:

$$\text{dom } \alpha_i \neq \text{dom } \beta_i \quad \text{or} \quad \text{ran } \alpha_i \neq \text{ran } \beta_i.$$

Since every subset with finite complement in $\mathbb{Z}$ is order isomorphic to $\mathbb{Z}$ we conclude that there exist monotone bijective maps $\gamma_i: \mathbb{Z} \rightarrow \text{dom } \alpha_i$ and $\delta_i: \text{ran } \alpha_i \rightarrow \mathbb{Z}$. Then we have that $\gamma_i \alpha_i \delta_i$ is an element of the group of units of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$, because $\text{dom}(\gamma_i \alpha_i \delta_i) = \text{ran}(\gamma_i \alpha_i \delta_i) = \mathbb{Z}$.

Suppose we have that $\text{dom } \alpha_i \neq \text{dom } \beta_i$. If there exists an integer $k \in \text{dom } \alpha_i$ such that $k \notin \text{dom } \beta_i$, then $(k)\gamma_i^{-1} \in \text{dom}(\gamma_i \alpha_i \delta_i)$ and $(k)\gamma_i^{-1} \notin \text{dom}(\gamma_i \beta_i \delta_i)$. If there exists an integer $k \in \text{dom } \beta_i$ such that $k \notin \text{dom } \alpha_i$, then $(k)\gamma_i^{-1} \in \text{dom}(\gamma_i \beta_i \delta_i)$ and $(k)\gamma_i^{-1} \notin \text{dom}(\gamma_i \alpha_i \delta_i)$. Therefore, we get that $\text{dom}(\gamma_i \beta_i \delta_i) \neq \text{dom}(\gamma_i \alpha_i \delta_i)$.

Suppose we have that $\text{ran } \alpha_i \neq \text{ran } \beta_i$. If there exists an integer $k \in \text{ran } \alpha_i$ such that $k \notin \text{ran } \beta_i$, then $(k)\delta_i \in \text{ran}(\gamma_i \alpha_i \delta_i)$ and $(k)\delta_i \notin \text{ran}(\gamma_i \beta_i \delta_i)$. If there exists an integer $k \in \text{ran } \beta_i$ such that $k \notin \text{ran } \alpha_i$, then $(k)\delta_i \in \text{ran}(\gamma_i \beta_i \delta_i)$ and $(k)\delta_i \notin \text{ran}(\gamma_i \alpha_i \delta_i)$. This implies that $\text{ran}(\gamma_i \beta_i \delta_i) \neq \text{ran}(\gamma_i \alpha_i \delta_i)$.

Since every translation on an arbitrary element of the group of units of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is a bijective map of the set of integers $\mathbb{Z}$, without loss of generality we can assume that the element $\gamma_i\alpha_i\delta_i$ is the unit of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$.

Next, we define elements $\gamma = (\gamma_1, \dots, \gamma_n)$ and $\delta = (\delta_1, \dots, \delta_n)$ of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ in the following way. For $i \in \mathbf{D}_{\alpha,\beta}$ we define γ_i and δ_i to be the elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ so constructed above. For $i \in \{1, \dots, n\} \setminus \mathbf{D}_{\alpha,\beta}$ we put γ_i and δ_i are the elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ such that $\gamma_i\alpha_i\delta_i = \gamma_i\beta_i\delta_i = \mathbb{I}_i$ is the unit of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$. The existence of so elements γ_i and δ_i in $\mathcal{SO}_\infty(\mathbb{Z})$ follows from Theorem 2.3 of [1] and the fact that the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is bisimple (see [4, Proposition 2.3]).

Hence we get that

$$\gamma\alpha\delta = \mathbb{I}, \quad \varpi = \gamma\beta\delta \neq \mathbb{I} \quad \text{and} \quad \varpi \mathfrak{C} \mathbb{I} \quad \text{in} \quad \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n).$$

Moreover, our construction implies that $\mathbf{D}_{\mathbb{I},\varpi} = \mathbf{D}_{\alpha,\beta}$. □

Lemma 2. *Let $\mathfrak{C}$ be a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Let α and β be two distinct $\mathfrak{C}$ -equivalent elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Then there exists an element ψ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbb{I}\mathfrak{C}\psi$, $\mathbf{D}_{\mathbb{I},\psi} = \mathbf{D}_{\alpha,\beta}$ and elements $\mathbb{I}$ and ψ are not $\mathcal{H}$ -equivalent in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$.*

Proof. If α and β are not $\mathcal{H}$ -equivalent elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$, then by case 2) of the proof of Lemma 1 we obtain that $\mathbb{I}\mathfrak{C}\varpi = \gamma\beta\delta$ and the elements $\mathbb{I}$ and ϖ are not $\mathcal{H}$ -equivalent in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$.

Next, we suppose that $\alpha \mathcal{H} \beta$ and put $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$. Then by Proposition 2.3 of [4] the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is bisimple and hence by Theorem 2.3 of [1] for every $i = 1, \dots, n$ there exist $\gamma_i, \delta_i \in \mathcal{SO}_\infty(\mathbb{Z})$ such that $\gamma_i\alpha_i\delta_i = \mathbb{I}_i$ is the unit of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ and $\gamma_i\beta_i\delta_i \neq \mathbb{I}_i$ for each $i \in \mathbf{D}_{\alpha,\beta}$. Since $\alpha \mathcal{H} \beta$ and by Proposition 2.3(v) of [2] the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is isomorphic to the direct power $(\mathcal{SO}_\infty(\mathbb{Z}))^n$ we conclude that $\gamma_i\beta_i\delta_i$ is an element of the group of units of $\mathcal{SO}_\infty(\mathbb{Z})$ for each $i \in \{1, \dots, n\}$, and moreover $\gamma_i\beta_i\delta_i = \mathbb{I}_i = \gamma_i\alpha_i\delta_i$ for any $i \in \{1, \dots, n\} \setminus \mathbf{D}_{\alpha,\beta}$.

We denote $\gamma = (\gamma_1, \dots, \gamma_n)$ and $\delta = (\delta_1, \dots, \delta_n)$ and put $\kappa = (\kappa_1, \dots, \kappa_n) = \gamma\beta\delta$. Then we have that $\mathbf{D}_{\alpha,\beta} = \mathbf{D}_{\mathbb{I},\kappa}$. Also the relation $\alpha \mathcal{H} \beta$ implies that $\mathbb{I} \mathcal{H} \kappa$, and since $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is an inverse semigroup we get that $\mathbb{I} \mathcal{H} \kappa^m$ for every integer m . By Proposition 2.2 of [4] the group of units of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is isomorphic to $\mathbb{Z}(+)$. Hence, this implies that without loss of generality we can assume that $(p)\kappa_i = p + m_i$, where $m_i \neq 0$, for every $i \in \mathbf{D}_{\alpha,\beta}$ because.

Next, for every integer $i = 1, \dots, n$ we define a partial map $\chi_i: \mathbb{Z} \rightarrow \mathbb{Z}$ in the following way:

- (a) if $i \in \{1, \dots, n\} \setminus \mathbf{D}_{\alpha,\beta}$, then we define $\chi_i: \mathbb{Z} \rightarrow \mathbb{Z}$ be the identity map;
- (b) if $i \in \mathbf{D}_{\alpha,\beta}$ and $m_i \geq 1$, then we define $\text{dom } \chi_i = \mathbb{Z}$, $\text{ran } \chi_i = \mathbb{Z} \setminus \{1, \dots, m_i\}$ and

$$(k)\chi_i = \begin{cases} k + m_i, & \text{if } k \geq 1; \\ k, & \text{if } k \leq 0; \end{cases}$$

- (c) if $i \in \mathbf{D}_{\alpha,\beta}$ and $m_i \leq -1$, then we define $\text{dom } \chi_i = \mathbb{Z}$, $\text{ran } \chi_i = \mathbb{Z} \setminus \{m_i, \dots, -1\}$ and

$$(k)\chi_i = \begin{cases} k, & \text{if } k \geq 0; \\ k + m_i, & \text{if } k \leq -1. \end{cases}$$

We put $\chi = (\chi_1, \dots, \chi_n)$. The definition of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ implies that χ and its inverse χ^{-1} are elements of $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Simple verifications show that $\mathbb{I} = \chi\chi^{-1} = \chi\mathbb{I}\chi^{-1}$. Also, since $\mathfrak{C}$ is a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ we conclude that $\mathbb{I} = \chi\mathbb{I}\chi^{-1}\mathfrak{C}\chi\kappa\chi^{-1}$.

Now simple calculations imply that

- (i) if $m_i > 0$ then

$$(k)\chi_i\kappa_i\chi_i^{-1} = \begin{cases} k + m_i, & \text{if } k \geq 1; \\ \text{undefined}, & \text{if } -m_i < k \leq 0; \\ k + m_i, & \text{if } k \leq -m_i, \end{cases}$$

and similarly

(ii) if $m_i < 0$ then

$$(k)\chi_i\kappa_i\chi_i^{-1} = \begin{cases} k + m_i, & \text{if } k \geq -m_i; \\ \text{undefined}, & \text{if } 0 \leq k < -m_i; \\ k + m_i, & \text{if } k \leq -1. \end{cases}$$

Next we put $\psi = \chi\kappa\chi^{-1}$, and hence we obtain that $\mathbb{I}\mathfrak{C}\psi$ but $\text{dom } \psi \neq \mathbb{Z}$. This completes the proof of our lemma. $\square$

Remark 3. The proof of Lemma 2 implies that for element $\psi = (\psi_1, \dots, \psi_n)$ the following property holds:

ψ_i is not $\mathcal{H}$ -equivalent to the unit of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ for every $i \in \mathbf{D}_{\alpha,\beta}$.

Proposition 4. Let $\mathfrak{C}$ be a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Let α and β be two distinct $\mathfrak{C}$ -equivalent elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Then there exists a non-unit idempotent ε in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbb{I}\mathfrak{C}\varepsilon$ and $\mathbf{D}_{\mathbb{I},\varepsilon} = \mathbf{D}_{\alpha,\beta}$.

Proof. Lemma 2 implies that there exists an element ψ of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\psi\mathbb{I}$, $\mathbf{D}_{\mathbb{I},\psi} = \mathbf{D}_{\alpha,\beta}$ and elements $\mathbb{I}$ and ψ are not $\mathcal{H}$ -equivalent in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Also, by Remark 3 for every integer $i \in \mathbf{D}_{\alpha,\beta}$ the element ψ_i is not $\mathcal{H}$ -equivalent to the unit $\mathbb{I}_i$ of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$. This implies that for every integer $i \in \mathbf{D}_{\alpha,\beta}$ at least one of the following conditions holds:

$$\psi_i\psi_i^{-1} \neq \mathbb{I}_i \quad \text{or} \quad \psi_i^{-1}\psi_i \neq \mathbb{I}_i \quad \text{in} \quad \mathcal{SO}_\infty(\mathbb{Z}).$$

Since $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is an inverse semigroup we have that $\mathbb{I}\mathfrak{C}\psi^{-1}$. This implies that $\mathbb{I}\mathfrak{C}\psi\psi^{-1}$ and $\mathbb{I}\mathfrak{C}\psi^{-1}\psi$, and hence we get that $\mathbb{I}\mathfrak{C}\varepsilon$, where $\varepsilon = \psi\psi^{-1}\psi^{-1}\psi$. The above arguments show that $\mathbf{D}_{\mathbb{I},\varepsilon} = \mathbf{D}_{\alpha,\beta}$. $\square$

Proposition 5. Let $\mathfrak{C}$ be a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Let α and β be two distinct $\mathfrak{C}$ -equivalent elements of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Then $\mathbb{I}\mathfrak{C}\varepsilon$ for any idempotent ε in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbf{D}_{\mathbb{I},\varepsilon} = \mathbf{D}_{\alpha,\beta}$.

Proof. By Proposition 4 there exists an idempotent ε of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbb{I}\mathfrak{C}\varepsilon$ and $\mathbf{D}_{\mathbb{I},\varepsilon} = \mathbf{D}_{\alpha,\beta}$. We fix an arbitrary non-unit idempotent $\iota \in \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\varepsilon \leq \iota$ in $E(\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n))$. Then we have that $\iota\mathbb{I} = \iota$ and hence the relation $\mathbb{I}\mathfrak{C}\varepsilon$ implies that $\iota = \iota\mathbb{I}\mathfrak{C}\varepsilon = \varepsilon\mathbb{I}$. Therefore, for every $i \in \mathbf{D}_{\alpha,\beta}$ there exists an idempotent ε_i° such that $\varepsilon_i^\circ\mathbb{I}$ and the set $\mathbb{Z} \setminus \text{dom } \varepsilon_i^\circ$ is singleton. We put $\{m_i\} = \mathbb{Z} \setminus \text{dom } \varepsilon_i^\circ$ for every integer $i \in \mathbf{D}_{\alpha,\beta}$. We fix an arbitrary integer p_i for $i \in \mathbf{D}_{\alpha,\beta}$ and define the map $\varrho_i: \mathbb{Z} \rightarrow \mathbb{Z}$ by the formula:

$$(j)\varrho_i = j - m_i + p_i, \quad \text{for every } j \in \mathbb{Z}.$$

Then ϱ_i is an element of the group of units of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ and hence $\varrho_i\varrho_i^{-1} = \varrho_i^{-1}\varrho_i = \mathbb{I}_i$ in $\mathcal{SO}_\infty(\mathbb{Z})$. Moreover, it is obvious that $\varrho_i^{-1}\varepsilon_i^\circ\varrho_i$ is an idempotent of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ such that $\text{dom}(\varrho_i^{-1}\varepsilon_i^\circ\varrho_i) = \mathbb{Z} \setminus \{p_i\}$. Also, we obtained that $\mathbb{I}_i = \varrho_i^{-1}\mathbb{I}_i\varrho_i\mathfrak{C}\varrho_i^{-1}\varepsilon_i^\circ\varrho_i$ in $\mathcal{SO}_\infty(\mathbb{Z})$. Now the definition of the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ implies that $\mathbb{I}\mathfrak{C}\iota_i^\circ$ for any idempotent ι in $\mathcal{SO}_\infty(\mathbb{Z})$, because every idempotent ι in the semigroup $\mathcal{SO}_\infty(\mathbb{Z})$ is equal to a product of finitely many idempotents of the form ι_i° , $i \in \{1, \dots, n\}$, with the property that the set $\mathbb{Z} \setminus \text{dom } \iota_i^\circ$ is singleton. Then for every idempotent ε of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ with the property $\mathbf{D}_{\mathbb{I},\varepsilon} = \mathbf{D}_{\alpha,\beta}$ we have that

$$\varepsilon = \varepsilon_{i_1}^\circ \cdot \dots \cdot \varepsilon_{i_k}^\circ, \quad \text{where } \{i_1, \dots, i_k\} = \mathbf{D}_{\alpha,\beta},$$

and hence $\mathbb{I}\mathfrak{C}\varepsilon$. This completes the proof of the proposition. $\square$

Theorem 6. Let $\mathfrak{C}$ be a congruence on the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. Then the following statements hold:

- (i) If $\Delta_{\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)} \subseteq \mathfrak{C} \subseteq \sigma_{[i_m]}$ for some $i_m \in \{1, \dots, n\}$, then either $\Delta_{\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)} = \mathfrak{C}$ or $\mathfrak{C} = \sigma_{[i_m]}$.
- (ii) If $\sigma_{[i_1, \dots, i_m]} \subseteq \mathfrak{C} \subseteq \sigma_{[i_1, \dots, i_m, i_{m+1}]}$, for any subset $\{i_1, \dots, i_m, i_{m+1}\} \subseteq \{1, \dots, n\}$, then either $\sigma_{[i_1, \dots, i_m]} = \mathfrak{C}$ or $\mathfrak{C} = \sigma_{[i_1, \dots, i_m, i_{m+1}]}$.

Proof. By Proposition 2.15 from [2] we have that for any collection $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ of distinct indices, $k \leq n$, and, hence, $\alpha\sigma_{[i_1, \dots, i_k]}\beta$ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ if and only if $\alpha\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ = \beta\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ$ for some idempotents $\varepsilon_{i_1}^\circ, \dots, \varepsilon_{i_k}^\circ \in \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$. This implies that $\mathbb{I}\sigma_{[i_1, \dots, i_k]}\varepsilon$ for every idempotent ε of the semigroup $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\mathbf{D}_{\mathbb{I}, \varepsilon} \subseteq \{i_1, \dots, i_k\}$. Then applying Proposition 4 we get the statement of the theorem. $\square$

For any proper subset of indices $I \subset \{1, \dots, n\}$ we define a map $\pi^I: \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n) \rightarrow \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ by the formula $(\alpha_1, \dots, \alpha_n)\pi^I = (\beta_1, \dots, \beta_n)$, where

$$\beta_i = \begin{cases} \alpha_i, & \text{if } i \in I; \\ \mathbb{I}_i, & \text{if } i \in \{1, \dots, n\} \setminus I. \end{cases}$$

Simple verifications show that such defined map $\pi^I: \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n) \rightarrow \mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ is a homomorphism. Let $\pi^{I^\#}$ be the congruence on $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ which is generated by the homomorphism π^I .

Proposition 7. *Let I be an arbitrary proper subset of $\{1, \dots, n\}$. Then $\pi^{I^\#}_{\min} = \sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]}$, where $\{i_1, \dots, i_k\} = \{1, \dots, n\} \setminus I$.*

Proof. Suppose that $\alpha(\sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]})\beta$ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ for some elements $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$. Proposition 2.15 of [2] implies that $\alpha\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ = \beta\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ$ for some idempotent $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ such that $\varepsilon_i = \mathbb{I}_i$ for all $i \in I$, i.e., $\alpha\varepsilon = \beta\varepsilon$. Then we have that $\alpha_i = \beta_i$ for all $i \in I$, and hence $\alpha\varepsilon^* = \beta\varepsilon^*$ for $\varepsilon^* = (\varepsilon_1^*, \dots, \varepsilon_n^*)$, where

$$\varepsilon_i^* = \begin{cases} \alpha_i^{-1}\alpha_i = \beta_i^{-1}\beta_i, & \text{if } i \in I; \\ \varepsilon_i, & \text{if } i \in \{1, \dots, n\} \setminus I. \end{cases}$$

It is obvious that $\varepsilon^*\pi^{I^\#}\alpha^{-1}\alpha\pi^{I^\#}\beta^{-1}\beta$. This implies the inclusion $\sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]} \subseteq \pi^{I^\#}_{\min}$.

Suppose that $\alpha\pi^{I^\#}_{\min}\beta$ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ for some elements $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\beta = (\beta_1, \dots, \beta_n)$. Then there exists an idempotent $\varepsilon = (\varepsilon_1, \dots, \varepsilon_n)$ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$ such that $\alpha\varepsilon = \beta\varepsilon$ and $\varepsilon\pi^{I^\#}\alpha^{-1}\alpha\pi^{I^\#}\beta^{-1}\beta$. The last two equalities imply that $\alpha_i^{-1}\alpha_i = \beta_i^{-1}\beta_i = \varepsilon_i$ for all $i \in I$. This and the equality $\alpha\varepsilon = \beta\varepsilon$ imply that $\alpha_i\varepsilon_i = \beta_i\varepsilon_i$ for all $i \in I$ and hence we obtain that $\alpha_i = \alpha_i\alpha_i^{-1}\alpha_i = \alpha_i\varepsilon_i = \beta_i\varepsilon_i = \beta_i\beta_i^{-1}\beta_i = \beta_i$ for all $i \in I$. Therefore we have that $\alpha\varepsilon^* = \beta\varepsilon^*$, where the idempotent $\varepsilon^* = (\varepsilon_1^*, \dots, \varepsilon_n^*)$ defined in the following way

$$\varepsilon_i^* = \begin{cases} \alpha_i^{-1}\alpha_i, & \text{if } i \in I; \\ \varepsilon_i, & \text{if } i \in \{1, \dots, n\} \setminus I. \end{cases}$$

This implies that $\alpha\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ = \beta\varepsilon_{i_1}^\circ \dots \varepsilon_{i_k}^\circ$. By Proposition 2.15 of [2] we get that $\alpha(\sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]})\beta$ in $\mathcal{SO}_\infty(\mathbb{Z}_{\text{lex}}^n)$, and hence we get that $\pi^{I^\#}_{\min} \subseteq \sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]}$. This completes the proof of equality $\pi^{I^\#}_{\min} = \sigma_{[i_1]} \circ \dots \circ \sigma_{[i_k]}$. $\square$

ACKNOWLEDGEMENTS

The authors are grateful to the referee for several useful comments and suggestions.

REFERENCES

- [1] A. H. Clifford and G. B. Preston, *The Algebraic Theory of Semigroups*, Vol. I., Amer. Math. Soc. Surveys 7, Providence, R.I., 1961; Vol. II., Amer. Math. Soc. Surveys 7, Providence, R.I., 1967.
- [2] O. Gutik and I. Pozdnyakova, *On monoids of monotone injective partial selfmaps of $L_n \times_{\text{lex}} \mathbb{Z}$ with co-finite domains and images*, Algebra Discr. Math. **17**:2 (2014), 256–279.
- [3] O. Gutik and D. Repovš, *Topological monoids of monotone, injective partial selfmaps of $\mathbb{N}$ having cofinite domain and image*, Stud. Sci. Math. Hungar. **48**:3 (2011), 342–353.
- [4] O. Gutik and D. Repovš, *On monoids of injective partial selfmaps of integers with cofinite domains and images*, Georgian Math. J. **19**:3 (2012), 511–532.
- [5] J. M. Howie, *Fundamentals of Semigroup Theory*, London Math. Monographs, New Ser. 12, Clarendon Press, Oxford, 1995.

- [6] M. Lawson, *Inverse Semigroups. The Theory of Partial Symmetries*, Singapore: World Scientific, 1998.
- [7] M. Petrich, *Inverse Semigroups*, John Wiley & Sons, New York, 1984.
- [8] V. V. Wagner, *Generalized groups*, Dokl. Akad. Nauk SSSR **84** (1952), 1119—1122 (in Russian).

FACULTY OF MECHANICS AND MATHEMATICS, NATIONAL UNIVERSITY OF LVIV, UNIVERSYTETSKA 1, LVIV, 79000,
UKRAINE

E-mail address: o_gutik@franko.lviv.ua, ovgutik@yahoo.com, pozdnyakova.inna@gmail.com